

FIG. 1. Sensitivity of the PCR assay. Shown are the results of PCR amplification of the serially diluted *L. donovani* (DD8) DNA analyzed on agarose gels. DNA was extracted from parasite cultures and amplified as described in Materials and Methods. Lane M, 1 kb Ladder (Gibco BRL); lane 1, 10 ng of DNA; lane 2, 1 ng of DNA; lane 3, 10 pg of DNA; lane 4, 1 pg of DNA; lane 5, 10 fg of DNA; lane 6, 1 fg of DNA.

Probe: Ld Ind kDNA

Human DNA: $\xrightarrow{100 \text{ ng}}$


Primer Set: Ldl1 & 2

Amt. Ld Ind DNA: 1 pg 0.1 pg 0.01 pg 0

(Kb)

0.87 —

0.6 —



1 2 3 4

FIG. 2. Sensitivity of PCR amplification of *Leishmania* kDNA followed by Southern blot analysis. The PCR contained 100 ng of human genomic DNA and the indicated amount of total DNA from *L. donovani* DD8. The PCR product was probed with parasite kDNA and exposed for about 1 h. Lane 4 represents a PCR containing only human DNA as a control.





FIG. 3. Amplification of parasite DNA from various strains and isolates of *Leishmania*. DNA (1 ng) isolated from parasite cultures was subjected to PCR and analyzed. Lane 1, *L. donovani* AG83; lane 2, *L. donovani* DD8; lane 3, *L. donovani* HCB8; lane 4, *L. donovani* CB6; lane 5, *L. donovani* HCB 7 (PKDL origin); lane 6, *L. donovani* 8; lane 7, *L. donovani* WR684; lane 8 *L. donovani* infantum; lane 9, *L. tropica* WR683; lane 10, *L. major* I.V. 39, lane M, 1-kb ladder, lane 11, *Plasmodium*; lane 12, *M. leprae*; lane 13, *M. tuberculosis*.

M 1 2 3 4 5 6 7 8 9 10 11



600 bp

FIG. 4. DNA amplification from recent field isolates of KA and KDL. DNA (1 ng) extracted from cultures of parasite isolates was used for PCR amplification. Lanes: M, 1-kb ladder; 1, KA-1; 2, KA-2; 3, KA-3; 4, KA-4; 5, KA-5; 6, PK-1; 7, PK-2; 8, PK-3; 9, PK-4; 10, PK-5; 11, isolate from a patient with cutaneous leishmaniasis.

U 013891-8




FIG. 5 PCR assay with clinical samples of KA and PKDL. DNA (100 ng) isolated from clinical samples was used for PCR amplification. Lane M, 1-kb ladder; lane 1, KA (bone marrow); lane 2, KA (blood); lane 3, malaria (blood); lane 4, tuberculosis (blood); lane 5, control from the area of endemicity (blood); lane 6, PKDL (skin lesion); lane 7, leprosy (lesion).

10086104.022702

1	gaatttcgccg	aaaastgacc	gaaaattgggc	caaaaaaccca	aactttttctg	gtccctccggg
61	tggggggcttc	ctcgcgaaaa	cggaaaaatgg	ctgcagaaat	cccgctcaaa	aaataagccaa
121	aaatgcacaa	aatcgcgcctc	gaggcggggaa	actggggggtt	gggtgaaaat	agggttcggg
181	ggaggggaaa	ctcgggggttc	ggacgtgtgt	ggatatggcc	tgggtgggga	ctttggagctt
241	ggctgtacct	gtatctgggtt	tggagcatctg	cttgggggtt	gggggttttgt	gtgggaaggg
301	ggctggcgct	atttcggagtg	acgttggcttc	ttttgataat	tgtatttctg	ttcaaatctg
361	attgggtctgg	ctcgacatac	gttgggtctgg	ctggaaattgg	atcgaggtctt	gaactctaac
421	gggggtcggg	gcttggactcg	gggttgaggga	gtttgtgggg	atagtcttgg	atgtatgatat
481	ggaaatcgagc	ttcctcttaac	ataaatatta	gttggggctg	ttgcattagt	ttgttcacg
541	ggagtagcct	cgaagcatctg	ggcgggagat	atcatattat	cggtagtata	atatcatatg
601	tatacggtat	agatatctta	taattgtagt	atattctaga	tctattctac	agtgataag
661	ctatgaactt	actagacata	atttgtattt	gatgctatag	tgctactact	agagtgagcc
721	ctatcactag	cttagcagttag	ctgaagcttc	ctaaatgggt	gggaactgggt	gtgaaggctg
781	gaacgaacac	tg				